



Best Agrolife Limited
(An ISO 9001 : 2015 Certified Company)

**Dual Mode Of Action -
Controls Multiple Stages
Of Fungus Development**



**Dose:
300 ml
Per Acre**



Presentation :

1 Ltr, 500 ml, 250 ml & 100 ml



ZOXIT (Azoxystrobin 11% + Tebuconazole 18.3% w/w SC)

Mode of Action ►

Zoxit is a suspensible concentrate fungicide for control of Fruit rot, Dieback and Powdery mildew in Chilli, Purple blotch (*Alternaria porri*) on Onion and Sheath blight on Rice Crop.

Recommendation ►

Crop (S)	Common Name of Diseases	Dosage / HA		Dilution in water (Litre)
		a.i. (gms)	Formulation (ml)	
Rice	Sheath blight (<i>Rhizoctonia solani</i>)	82.5+137.25	750	800
Onion	Purple blotch (<i>Alternaria porri</i>)	82.5+137.25	750	800
Chilli	Fruit rot, Powdery mildew, Dieback	72+120	600	500-750
Potato	Early blight (<i>Alternaria solani</i>), Late blight (<i>Phytophthora infestans</i>)	82.5+137.25	750	500
Wheat	Yellow rust (<i>Puccinia striiformis</i>)	82.5+137.25	750	500
Tomato	Early blight (<i>Alternaria solani</i>)	82.5+137.25	750	500
Grape	Downy mildew, Powdery mildew	82.5+137.25	750	500
Apple	Scab (<i>Venturia inaequalis</i>), Powdery mildew (<i>Podosphra leucotncha</i>), Premature leaf fall (<i>Marssonina coronaria</i>)	0.11+0.183/L	1.0/L of water	8-12 L/ tree



Method of Use ►

Take the measured quantity of fungicide and add to it a small quantity of water. Stir thoroughly using a stick or rod and then add remaining quantity of water, stir well to form final spray solution.

Time of Application ►

To be applied preventively and to be repeated at 15 days interval depending upon prevailing weather conditions and disease pressure.