




Best Agrolife Limited

(An ISO 9001 : 2015 Certified Company)

The Duty To Gun Down Cyperus Weeds



Dose:
36 g
Per Acre

Presentation : 36 g & 3.6 g



HEADSHOT (Halosulfuron Methyl 75% WG)



Mode of Action ►

Headshot (Halosulfuron Methyl 75% WG) is a post emergence broad spectrum herbicide that belongs to the chemical group of sulfonylurea compound. Halosulfuron Methyl 75% WG has excellent foliar application efficacy against *Cyperus rotundus* in Sugarcane crop, *Cyperus rotundus* and *Cyperus iria* in Maize and Bottle Gourd crops.

It is taken up through the roots and shoots via the xylem and downwards via the phloem. It accumulates in the plant and inhibits the enzyme aceto-lactate synthate (ALS), which is responsible for purchasing essential amino acids. The affected weeds cannot, therefore, maintain proteins for new cells. The whole process is driven by the plant's use of water (transpiration).

- **Low-Dose Efficacy**
- **Excellent Control**
- **Multiple Features**
- **Improved Healthier Crop & Yield**



Recommendation ►

Crop (S)	Common Name of Weeds	Dosage / HA		Dilution in water (Litre)
		ai (gm)	Formulation (ml)	
Sugarcane	<i>Cyperus rotundus</i> L.	60-67.5	80-90	375
Maize	<i>Cyperus rotundus</i> L., <i>Cyperus iria</i> L.	67.5	90	375
Bottle Gourd	<i>Cyperus rotundus</i> L., <i>Cyperus iria</i> L.	67.5	90	375

Method of Use ►

Slowly add the required quantity of herbicide to water and stir well with a stick or rod till it is properly emulsified.

Time of Application ►

Crop Stage for application: Vegetative phase.

HEADSHOT