



**Best Agrolife Limited**

(An ISO 9001 : 2015 Certified Company)

▶ Quick ▶ Safe ▶ Sure



Dose:  
**2.5 ml**  
Per Ltr. Water

Presentation :  
1Ltr, 500 ml, 250 ml & 100 ml

<40













# PROPIQUE (Propaquizafop 10% EC)



## Mode of Action ▶

Propique (Propaquizafop 10% EC) is a post emergence herbicide predominantly effective against monocot weeds in Soybean, Black gram and Onion.

 	 	 	 	 
Creating new impetus With advanced crop Solutions	A powerful shield for abundant & healthy harvest produce	The power of outstanding control on unwanted annual and perennial grasses	quickly absorbed by the leaves and translocated from the foliage	Prompt action & improved systemic strength

## Recommendation ▶

Crop (S)	Common Name of Weeds	Dosage / HA		Dilution in water (Litre)
		ai (gm)	Formulation (ml)	
Soybean	Echinochloa coloum, Echinochloa crusgalli, Digitaria sanguinalis Dactyloctenium aegyptium Eleusine indica	50- 75 g	500- 750 ml	500-750
Onion	Digitaria sanguinalis Dactyloctenium aegyptium Echinochloa coloum Phalaris minor	62.5 g	625 ml	500
Black Gram	Echinochloa coloum, Digitaria sanguinalis Dactyloctenium aegyptium Eleusine indica	75- 100 g	750- 1000 ml	500-750

## Method of Use ▶

Shake the bottle before use measure out requisite quantity of the herbicide and mix it well with small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable sprayer.

**Crop Stage for application:** When weeds are in 2-4 leaf stage. **Application Technique:** Foliar application using high volum sprayers and ASPEE-HTP power sprayer fitted with flat fan or flood jet nozzle. **Equipment Type:** knapsack sprayer with flat fan/flood zet nozzle. **NozzleType:** Flat fan/flood zet nozzle.

## Time of Application ▶

**Soybean/Black gram :** 15 - 20 days after sowing, **Onion:** 15 days after transplanting, **Tomato / Chilli :** 20 days after transplanting. **Application Timing:** Preferably in the morning and evening hours.

PROPIQUE