



Best Agrolife Limited

(An ISO 9001 : 2015 Certified Company)

Targets


Broad Leaf Weeds
As Well As Sedges
With Wide Selectivity



Dose:
15 g
Per Acre

Presentation : 15 g

MAJOUR (Chlorimuron Ethyl 25% WP)




Mode of Action ►

Majour is an early post-emergence systemic herbicide belonging to the chemical family of Sulfonyl Urea herbicide. It controls broad leaf weeds and sedges with a wide selectivity and broad spectrum activity in Soybean and Transplanted Rice crops.

Recommendation ►

Crop (S)	Common Name of Weeds	Dosage / HA		Dilution in water (Litre)
		ai (gm)	Formulation (gm)	
Rice (Transplanted)	Echinochloa crusgalli, Eclipta alba, Commelina benghalensis, Chenopodium album, Cyperus rotundus	6	24	500-600
Soybean	Phyllanthus niruri, Celosia argentia, Cyperus rotundus, Digeria arvensis, Cucumis trigonus, Cyperus iria, Parthenium hysterophorus, Acalypha indica, Trianthema portulacastrum, Commelina benghalensis, Caesulia axillaris	9	36	300



Method of Use ►

Plant Protection Equipment: Knapsack sprayer fitted with flat fan/ flood jet nozzle.

Mode of Application: As spray suspension in 500-600 litre water per hectare.

Time of Application ►

Soybean : 10-20 days after sowing, (once)

Rice: 5-10 days after transplanting, (once)